

Nonparametric and Distribution- Free Statistics



Odisha Social Science
Research & Consultancy

CONTENTS

- **Introduction**
- **The Sign Test**
- **The Wilcoxon Signed- Rank Test for Location**
- **The Median Test**
- **The Mann-Whitney Test**
- **The Kolmogorov- Smirnov Goodness-of-fit Test**
- **The Kruskal-Wallis one-way analysis of variance by ranks**

Introduction

- Those procedures that test hypotheses that are not statements about population parameters are classified as *nonparametric*.
- Those procedures that make no assumption about the sampled population are called *distribution free* procedures.
- However, both these terms Nonparametric and distribution free are used interchangeably and under the *common heading 'Nonparametric'*.

Advantages:

- **Allow for the testing of hypothesis that are not statements about population parameter values.**
- **It may be used when the form of the sampled population is unknown.**
- **Computationally easier.**
- **May be applied when the data being analyzed consists merely of rankings or classifications.**

Disadvantages:

- **Application of Nonparametric procedure with data that can be handled with a parametric procedure results in a waste of data.**
- **Application of the Nonparametric tests may be laborious for large samples.**

The Sign Test

- The 't' test is not strictly valid for testing hypothesis regarding single population mean or mean difference between pairs of observations unless the relevant populations are at least approximately normally distributed.
- A frequently used nonparametric test that doesn't depend on the assumptions of 't' test is the sign test.
- This test focuses on the median rather than the mean as a measure of central tendency or location.
- The only assumption underlying the test is that the distribution of the variable of the interest is continuous.

Example: In a school for the mentally retarded, 10 girls selected at random received special instruction in personal care and grooming. Two weeks after completion of the course of instruction the girls were interviewed by a nurse and a social worker who assigned each girl a score based on her general appearance. The investigators believed that the score achieved the level of an ordinal scale. The scores are shown in the table. We wish to know if we can conclude that the median score of the population from which we assume this sample to have been drawn is different from 5

Table: General appearance Scores of 10 mentally Retarded Girls

Girl	Score	Girl	Score	Girl	Score
1	4	5	9	9	6
2	5	6	6	10	6
3	8	7	10		
4	8	8	7		

Data: The data are given in the table

Assumption: We assume that the measurements are taken on a continuous variable.

Hypothesis: H_0 : The population median is 5

H : The population median is not 5

Let $\alpha=0.05$

Test Statistic (TS): This will depend upon the H_A :

If $H_A: P(+) > P(-)$ A sufficiently smaller number of – signs causes rejection of H_0 . **TS** = ‘K’ = No. of – signs.

If $H_A: P(+) < P(-)$ A sufficiently smaller number of + signs causes rejection of H_0 . **TS** = ‘K’ = No. of + signs.

If $H_A: P(+) \neq P(-)$ Either a sufficiently smaller number of + signs or – signs causes rejection of H_0 . **TS** = ‘K’ Less frequently occurring signs of the two. This is the TS in the present example.

Distribution of Test statistic: If we assign a +sign to those score that lies above the hypothesized median of 5 and a – sign to those that fall below. Then the table is

Girl	1	2	3	4	5	6	7	8	9	10
Score	-	0	+	+	+	+	+	+	+	+

From the table we constitute a set of n independent random variable from the Bernoulli population with parameter p. If we let $K =$ the TS, the sampling distribution of K is the binomial distribution with parameter $p=0.5$ if H_0 is true.

Decision Rule:

For Reject H_0 if the ‘p’ value for the computed test statistic is less than or equal to 0.05.

Calculation of test statistic: Determine the probability of observing S or fewer –signs when given a sample of size n and parameter $p=0.5$.

Decision rule: The decision rule depend on the alternative hypothesis.

For $H_A: P(+) \neq P(-)$ reject H_0 if (given H_0 is true) the probability of obtaining a value of 'S' as extreme as or more extreme than the actually computed is equal to or less than $\alpha/2$.

$$P(k \leq x | n, p) = \sum_n C_k p^k q^{n-k}$$

For the example

$$\begin{aligned} & {}_9C_0(0.5)^0(0.5)^{9-0} + {}_9C_1(0.5)^1(0.5)^{9-1} \\ &= .00195 + .01758 \\ &= .0195 \end{aligned}$$

Here there are fewer minuses so we focus our attention on minuses rather than pluses.

Statistical decision: If the number of minuses is so small that the probability of observing this few of fewer is less than 0.025, we will reject the null hypothesis. The probability .0915 is less than .025 so reject the null hypothesis.

Conclusion: The median score is not 5.

p value: The p value for this test is $2(.0195) = 0.0390$

Sign Test-Paired Data:

Conditions

- The data to be analyzed consist of observation in matched pairs
- The measurement scale is weak.
- Sign test may be employed to test the null hypothesis that the median difference is zero

- Say (Y_i, X_i) are pairs of matched scores, $i = 1, 2, \dots, n$

One of the matched scores, say Y_i is subtracted from the other score X_i . If $Y_i < X_i$, the sign of the difference is + and if $Y_i > X_i$ the sign of the difference is - Then the null hypothesis is

$H_0: P(+) = P(-) = 0.5$ or $P(X_i > Y_i) = P(X_i < Y_i) = 0.5$

$H_A: P(+) \neq P(-)$ or $H_A: P(+) > P(-)$ or $H_A: P(+) < P(-)$

Example:

A dental research team wished to know if teaching people how to brush their teeth would be beneficial. 12 pairs of patients seen in a dental clinic were obtained by carefully matching on such factors as age, sex, intelligence and initial oral hygienic scores. One member of each pair received instruction on how to brush the teeth and other oral hygienic matters. Six months later all 24 subjects were examined and assigned an oral hygienic score by a dental hygienist unaware of which subjects had received the instruction. A low score indicates a high level of oral hygienic. The results are shown in Table.

Pair No	1	2	3	4	5	6	7	8	9	10	11	12
Inst. (Xi)	1.5	2.0	3.5	3.0	3.5	2.5	2.0	1.5	1.5	2.0	3.0	2.0
Not inst. (Yi)	2.0	2.0	4.0	2.5	4.0	3.0	3.5	3.0	2.5	2.5	2.5	2.5

Solution:

Assumption: We assume that the population of difference between two pairs of scores is a continuous variable.

Hypothesis:

H_0 : The median of the differences is zero [$P(+)\geq P(-)$]

H_A : The median of the differences is negative [$P(+)<P(-)$]

Let $\alpha=0.05$.

Test Statistic: The test statistic is the number of +signs.

Distribution of Test Statistic: The sampling distribution of k is the binomial distribution with parameters n and 0.5 if H_0 is true.

Decision rule: Reject H_0 if $P(S \leq 2 | 11, 5) \leq 0.05$.

Calculation Of Test Statistic: The score difference have been obtained for each pair. Then the result is in Table.

Pair	1	2	3	4	5	6	7	8	9	10	11	12
Score	-	0	-	+	-	-	-	-	-	-	+	-

The nature of the hypothesis indicates a one sided test so that all of $\alpha = 0.05$ is associated with the rejection region, which consists of all values of S (equal to number of +signs).

The experiment yielded 1zero , 2pluses and 9 minuses.

When we eliminate 0, the effective sample size is $n=11$ with 2 pluses and 9minuses. Since a small no of +signs will cause rejection of the null hypothesis, test statistic is $S=2$.

Statistical Decision: The probability of obtaining no more than 2 pluses out of eleven tries when the null hypothesis is true is

$P(S \leq 2 | 11, 5) = \sum_{11} C_S (.5)^S (.5)^{11-S}$. This probability is to be .0327. since .0327 is less than .05, we must reject H_0 .

Conclusion: Then we conclude that the median difference is negative and the instruction was beneficial.

p value: The p value is 0.0327

Sample size

- When the sample size is large and p is close to 0.5 the binomial distribution may be approximated by normal distribution.
- According to the rule of thumb when both np and nq are greater than 5, normal approximation is appropriate.
- Under H_0 $p = 0.5$, a sample size of 12 would satisfy normal approximation.
- In this situation normal z test can be applied for sign test.
- The test statistic here

$$z = \frac{(k \pm 0.5) - 0.5n}{0.5 \sqrt{n}}$$

The Wilcoxon Signed- Rank Test for Location

- To test a null hypothesis about a population mean , but for some reason neither z nor t is an appropriate test statistic.
- Wilcoxon Signed- Rank Test is a appropriate test which makes use of the magnitudes of the differences between measurements and a hypothesized location parameter rather than just the signs of the differences.

Assumptions:

- The sample is random
- The variable is continuous.
- The population is symmetrically distributed about its mean μ .
- The measurement scale is at least interval.

Hypothesis:

The hypothesis are :

(a) $H_0: \mu = \mu_0$

$H_A: \mu \neq \mu_0$

(b) $H_0: \mu \geq \mu_0$

$H_A: \mu < \mu_0$

(c) $H_0: \mu \leq \mu_0$

$H_A: \mu > \mu_0$

Calculation:

1. Subtract the hypothesized mean μ_0 from each observation x_i , to obtain $d_i = x_i - \mu_0$.
2. Rank the usable d_i from the smallest to the largest without regard to the sign of d_i . If two or more $|d_i|$ are equal, assign each tied value the mean of the rank positions the tied values occupy.
3. Assign each rank the sign of the d_i that yields that rank.
4. Find T_+ , the sum of the ranks with +ve sign and T_- , the sum of the ranks with -ve sign.

The Test Statistic:

The Wilcoxon test statistic is either T_+ or T_- , depending on the nature of the alternative hypothesis.

- When the alternative hypothesis is two-sided ($\mu \neq \mu_0$) either a sufficiently small value of T_+ or a sufficiently small value of T_- will cause us to reject $H_0: \mu = \mu_0$. The test statistic then is T_+ or T_- , whichever is smaller.
- When $H_0: \mu \geq \mu_0$ is true we expect our sample to yield a large value of T_+ . Therefore, when the one-sided alternative hypothesis state that the true population mean is less than the hypothesized mean ($\mu < \mu_0$), a sufficiently small value of T_+ will cause rejection of H_0 and T_+ is the test statistic.
- When $H_0: \mu \leq \mu_0$ is true we expect our sample to yield a large value of T_- . Therefore, for the one-sided alternative $H_A: \mu > \mu_0$, a sufficiently small value of T_- will cause rejection of H_0 and T_- is the test statistic.

Critical Value:

The following are the decision rules for the three possible alternative hypothesis:

- (a) $H_A: \mu \neq \mu_0$. Reject H_0 at the α level of significance if the calculated T is smaller than or equal to the tabulated T for n and preselected $\alpha/2$. Alternatively, we may enter Table K with n and our calculated T is less than or equal to our Stated level of significance. If so, we may reject H_0 .
- (b) $H_A: \mu < \mu_0$. Reject H_0 at the α level of significance if the calculated T_+ is less than or equal to the tabulated T for n and preselected α .
- (c) $H_A: \mu > \mu_0$. Reject H_0 at the α level of significance if the calculated T_- is less than or equal to the tabulated T for n and preselected α .

Example:

Cardiac output(lit/min) was measured by thermodilution in a simple random sample of 15 postcardiac surgical patients in the left lateral position. The results were

4.91	4.10	6.74	7.27	7.42	7.50	6.56	4.64
5.98	3.14	3.23	5.80	6.17	5.39	5.77	

We wish to know if we can conclude on the basis of these data that the population mean is different from 5.05.

Solution:

Assumption: We assume that the requirements for the application of the Wilcoxon signed-ranks test are met.

Hypothesis: $H_0: \mu = 5.05$

$H_A: \mu \neq 5.05$ and Let $\alpha = 0.05$

Test Statistic: The test statistic will be T_+ or T_- , whichever is smaller.

Distribution of Test Statistic: Critical values of the test statistics are given in Table K.

Decision Rule: The decision rule is reject H_0 if the computed value of T is less than or equal to 25, the critical value for $n=15$, and $\alpha/2 = .0240$, the closest value to .0250 in Table K.

Calculation of test statistic: The calculation of the test statistic is in the table-

Table: Calculation of the test statistic

Cardiac output	$d_i = x_i - 5.05$	Rank of $ d_i $	Signed rank of $ d_i $
4.91	-.14	1	-1
4.10	-.95	7	-7
6.74	+1.69	10	+10
7.27	+2.22	13	+13
7.42	+2.37	14	+14
7.50	+2.45	15	+15
6.56	+1.52	9	+9
4.64	-.41	3	-3
5.98	+.93	6	+6
3.14	-1.91	12	-12
3.23	-1.82	11	-11
5.80	+.75	5	+5
6.17	+1.12	8	+8
5.39	+.34	2	+2
5.77	+.72	4	+4

$T_+ = 86$, $T_- = 34$, $T = 34$.

Statistical Decision: Since 34 is greater than 25, we are unable to reject H_0 .

Conclusion: We conclude that the population mean may be 5.05

p value: From Table K we see that the p value is $p = 2(.0757) = 0.1514$.

The Median Test

- A nonparametric procedure that may be used to test the null hypothesis that two independent samples have been drawn from populations with equal medians is the median test.

Example:

Do urban and rural male junior high school students differ with respect to their level of mental health?

Urban	Rural	Urban	Rural
35	29	25	50
26	50	27	37
27	43	45	34
21	22	46	31
27	42	33	
38	47	26	
23	42	46	
25	32	41	

Solution:

Assumption: The assumptions are the samples are selected independently ,at random from their respective population and the populations are of the same form, differing only in location and the variable of interest is continuous .

Hypothesis: $H_0: M_U = M_R$ and $H_A: M_U \neq M_R$.

M_U is the median score of the sampled population of urban students and M_R is the median score of the sampled population of rural students.

Let $\alpha = 0.05$.

Test Statistic: The test statistic is X^2 as computed.

Distribution of test statistic: When H_0 is true and the assumptions are met, X^2 with 1 degree of freedom.

Decision Rule: The decision rule is reject H_0 if the computed value of X^2 is ≥ 3.841 (since $\alpha=0.05$)

Calculation of test statistic: The first step in calculating the test statistic is to compute the common median i.e. $(33+34)/2 = 33.5$. The result according to for each group the number of observations falling above and below the common median is

	Urban	Rural	Total
No. of scores above median	6	8	14
No. of scores below median	10	4	14
Total	16	12	28

The test statistic is $\chi^2 = \frac{28[(6)(4) - (8)(10)]^2}{(16)(12)(14)(14)} = 2.33$

Statistical decision: Since $2.33 < 3.841$, the critical value of X^2 with $\alpha=.05$ and 1 degree of freedom, we are unable to reject the null hypothesis on the basis of these data.

Conclusion: The two samples may have been drawn from populations with equal median.

p value: The p value is $p > 0.10$ since $2.33 < 2.706$.

Handling Values Equal to the Median:

When one or more observed values will be exactly equal to the common median and, hence, will fall neither above nor below it and if $n_1 + n_2$ is odd, at least one value will always be exactly equal to the median one solution is drop them from the analysis if $n_1 + n_2$ is large and there are only a few values that fall at the combined median or we may dichotomize the scores into those that exceed the median and those that do not, in which case the observations that equal to the median will be counted in the second category.

Median Test extension:

The median test extends logically to the case where it is desired to test the null hypothesis that $k \geq 3$ samples are from populations with equal medians. For this test a $2 \times k$ contingency table may be constructed by using the frequencies that fall above and below the median computed from combined samples. If conditions as to sample size and expected frequencies are met, χ^2 may be computed and compared with the critical χ^2 with $k-1$ degrees of freedom.

The Mann-Whitney Test

- If for testing the desired hypothesis, there is available a procedure that makes use of more of the information inherent in the data, that procedure should be used if possible
- Such a nonparametric procedure that can often be used instead of the median test is called the Mann-Whitney Test.

Assumption:

- The two sample of size n and m , respectively available for analysis have been independently and randomly drawn from their respective populations.
- measurement scale is at least ordinal.
- variable of interest is continuous.
- If the population differ at all, they differ only with respect to their medians.

Hypothesis:

When these assumptions are met we may test the null hypothesis that the two populations have equal medians against either of the three possible alternatives:

- (1) the populations don't have equal medians(two sided test)**
- (2) the median of population 1 is larger than the median of population 2(one sided test)**
- (3) the median of population 1 is smaller than the median of population 2(one-sided test).**

Example:

A researcher designed an experiment to assess the effects of prolonged inhalation of cadmium oxide. 15 laboratory animals served as experimental subjects, while 10 similar animals served as controls. The variable of interest was hemoglobin level following the experiment. If we can conclude that prolonged inhalation of cadmium oxide reduces hemoglobin level

Solution:

Assumption: We presume that the assumptions of the Mann-Whitney test are met.

Table: Hemoglobin determination (grams) for 25 laboratory animals

Exposed animals (X)	Unexposed animals (Y)
14.4	17.4
14.2	16.2
13.4	17.1
16.5	17.5
14.1	15.0
16.6	16.0
15.9	16.9
15.6	15.0
14.1	16.3
15.3	16.8
15.7	
16.7	
13.7	
15.3	
14.0	

The null and alternative hypothesis are as follows:

$$H_0: M_X \geq M_Y$$

$H_A: M_X < M_Y$ where M_X is the median of a population of animals exposed to cadmium oxide and M_Y is the median of a population of animals not exposed to the substance. Let $\alpha = 0.05$.

Test statistic :

- 1. Two samples are combined**
- 2. All observations are ranked from smallest to largest keeping track of the sample to which each observations belongs.**
- 3. Tied observations are assigned a rank equal to the mean of the rank position for which they are tied.**

$$\text{The test statistic: } T = S - \frac{n(n+1)}{2}$$

Where n is the size of sample X and S is corresponding sum of ranks.

Distribution Of test: Critical values from the distribution of the test statistic are given in Table L for various levels of α .

Decision Rule: In general, for one-sided tests of the type illustrated here the decision rule is :

Reject $H_0: M_X \geq M_Y$ if the computed t is less than $w\alpha$ is the critical value of T obtained by entering Table L with n , the number of X observations, m , the number of Y observations, and α , the chosen level of significance.

Calculation of test statistic: $S=145$, so that

$$T = 145 - \frac{15(15+1)}{2} = 25$$

Statistical decision:

when we enter Table L with $n=15$, $m=10$, and $\alpha = 0.05$, we find the critical value of w_α to be 45. since $25 < 45$, we reject H_0 .

Conclusion: We conclude that M_x is smaller than M_y .
This leads to the conclusion that prolonged inhalation of cadmium oxide does reduce the hemoglobin level.

p value: since $22 < 25 < 30$, we have for this test $0.005 > p > 0.001$.

The Kolmogorov- Smirnov Goodness-of-fit Test

- The Kolmogorov's work concerned with the one-sample case as discussed and Smirnov's work deals with the case involving two samples in which interest centers on testing the hypothesis that the distributions of the two parent populations are identical .

Test Statistic:

- Kolmogorov- Smirnov goodness-of-fit test a comparison between some theoretical cumulative distribution function, $F_T(x)$, and a sample cumulative distribution function $F_S(x)$.
- The difference between the theoretical cumulative distribution function, $F_T(x)$, and the sample cumulative distribution function $F_S(x)$, is measured by the statistic D which is the greatest vertical distance between $F_S(x)$ and $F_T(x)$.

- When a two sided test is appropriate, that is when the hypothesis are

$H_0: F(x) = F_T(x)$ for all x from $-\infty$ to $+\infty$

$H_A: F(x) \neq F_T(x)$ for at least one x the statistic is

$$D = \sup_x |F_s(x) - F_t(x)|$$

which is read, “d equals the supremum (greatest) over all x , of the absolute value of the difference $F_s(x) - F_t(x)$.”

- The null hypothesis is rejected at the α level of significance if the computed value of D exceed the for $1 - \alpha$ (two sided) and the sample size n .

Assumption:

The assumptions are

1. Sample is a random sample
2. Hypothesized distribution $F_T(x)$ is continuous.

- When values of D are based on a discrete theoretical distribution, the test is conservative.
- When the test is used with discrete data, then, the true probability of committing a type-I error is at most equal to α , the stated level of significance.

Example:

Fasting, blood glucose determinations made on 36 nonobese, apparently healthy, adult males in table. We have to test if we may conclude that these data are not from a normally distributed population with a mean of 80 and a standard deviation of 6.

Solution:

Assumption: We assume that the sample available is a simple random sample from a continuous population distribution.

Hypothesis: $H_0: F(x) = F_T(x)$ for all x from $-\infty$ to $+\infty$

$H_1: F(x) \neq F_T(x)$ for at least one x and Let $\alpha=0.05$

Table: Fasting blood glucose values(mg/100ml) for 36 Nonobese, apparently healthy, adult males

75	92	80	80	84	72
84	77	81	77	75	81
80	92	72	77	78	76
77	86	77	92	80	78
68	78	92	68	80	81
87	76	80	87	77	86

Distribution of test statistic: Critical value of the test statistic for selected values of α are given in Table M.

Decision rule: Reject H_0 if the computed value of D exceeds 0.221, the critical value of D for $n=36$ and $\alpha=0.05$.

Calculation of test statistic: First step is to compute values of $F_S(x)$. Each value of $F_S(x)$ is obtained by dividing the corresponding cumulative frequency by the sample size like first value of $F_S(x) = 2/36 = 0.0556$

Value of $F_T(x)$ can be obtained by converting each observed value of x to a value of the standard normal variable, z . Then find the area between z and z_0 .

Table: Value of $F_S(x)$

X	F	C.F	$F_S(x)$
68	2	2	.0556
72	2	4	.1111
75	2	6	.1667
76	2	8	.2222
77	6	14	.3889
78	3	17	.4722
80	6	23	.6389
81	3	26	.7222
84	2	28	.7778
86	2	30	.8333
87	2	32	.8889
92	4	36	1.000

Table: Values of $F_T(x)$

X	$Z=(x-80)/6$	$F_T(x)$
68	-2.00	.0228
72	-1.33	.0918
75	-.83	.2033
76	-.67	.2514
77	-.50	.3085
78	-.33	.3707
80	.00	.5000
81	.17	.5675
84	.67	.7486
86	1.00	.8413
87	1.17	.8790
92	2.00	.9772

Statistical decision: Reference to Table M reveals that a computed D of 0.1547 is not significant at any reasonable level. So we are not willing to reject H_0 .

Conclusion: The sample may have come from the specified distribution.

The Kruskal-Wallis one-way analysis of variance by ranks

- When the populations from which the samples are drawn are not normally distributed with equal variances and when the data for analysis consists only of ranks, a nonparametric alternative to the one way analysis of variance may be used to test the hypothesis of equal location parameter.
- The best known of these procedures is the Kruskal-Wallis one way analysis of variance by ranks.
- The application of the test involves the steps
 - The $n_1, n_2, \dots, n_k$ observations from the k samples are combined in to a single series of size n and arranged in order of magnitude from smaller to largest.
 - Then the observations are then replaced by ranks from 1 to n from smallest to the largest value.

➤ The ranks assigned to observations in each of the k groups are added separately to give k rank sums.

➤ The test statistic

$$H = \frac{12}{n(n+1)} \sum \frac{R_j^2}{n_j} - 3(n+1)$$

Is computed.

k = the number of samples

n_j = the number of observation in the j th sample

n = the number of observations in all samples combined

R_j = the sum of the ranks in the j th sample.

➤ When there are 3 samples and five or fewer observations in each sample, the significance of the computed H is determined by consulting Table.

➤ When there are more than five observations in one or more of the samples, H is compared with tabulated values of χ^2 with $k-1$ degrees of freedom.

Example:

The effects of two drugs on reaction time to a certain stimulus were studied in three samples of experimental animals. Sample-III served as a control while the animals in sample-I were treated with drug A and those in sample II were treated with drug B prior to the application of the stimulus. Table shows the reaction times in seconds of the animals.

Can we conclude that the three populations represented by the three samples differ with respect to reaction time? We can so conclude if we can reject the null hypothesis that the three populations do not differ in their reaction times.

Table:1 Reaction time in seconds of 13 experimental animal

Sample		
I	II	III
17	8	2
20	7	5
40	9	4
31	8	3
35		

Solution:

Assumption: The samples are independent random samples from their respective populations. The measurement scale employed is at least ordinal. The distribution of the values in the sampled population are identical except for the possibility that one or more of the populations are composed of values that tend to be larger than the of the other populations.

Hypothesis:

H_0 : The population centers are equal

H_A : At least one of the populations tends to exhibit larger values than at least one of the other populations.

Let $\alpha=0.01$

Distribution of test statistic: critical values of H for various sample sizes and α levels are given in Table N.

Decision rule: The null hypothesis will be rejected if the computed value of H is very large that the probability of obtaining a value that large or larger when H_0 is true or equal or less than the chosen significance level, α .

Calculation of test statistic: When the three samples are combined in to a single series and ranked , the Table of ranks shown in Table 2.

Table:2 The data of Table 1 replaced by Ranks

Sample		
I	II	III
9	6.5	1
10	5	4
13	8	3
11	6.5	2
12		
$R_1=55$	$R_2=26$	$R_3=10$

From the Table data and equation we obtain

$$H = \frac{12}{13(13+1)} \left[\frac{(55)^2}{5} + \frac{(26)^2}{4} + \frac{(10)^2}{4} \right] - 3(3+1)$$
$$= 10.68$$

Statistical decision: Table N shows that when the n_j are 5, 4 and 4, the probability of obtaining a value of $H \geq 10.68$ is less than 0.009. The null hypothesis can be rejected at the .01 level of significance.

Conclusion: We conclude that there is a difference in the average reaction time among the three populations.

p value: $p < 0.009$.